

Frascati Physics Series Vol. XXX (1997), pp. 000-000
 CONFERENCE TITLE - Conference Town, Oct 3rd, 1996

NEW RESULTS FROM VES.

Valeri Dorofeev, VES collaboration
IHEP, Protvino, Russia, RU-142284

ABSTRACT

The results of the partial wave analysis(PWA) of the $\pi^+\pi^-\pi^-$ and $\omega\pi^-\pi^0$ systems are presented. The a_3 and $a_4(2040)$ signals are observed in the $\rho(770)\pi$ and $f_2(1270)\pi$ channels. Indications of the a'_1 meson existence was found in the 1^+0^+ $\rho\pi$ S-wave. The decay branching ratio of the $a_2(1320)^-$ to $\omega\pi^-\pi^0$ was measured. The 2^+1^+ wave shows a broad bump at $M \approx 1.7 GeV$. The decays of the $\pi_2(1670)$, $a_4(2040)$ and $\pi(1740)$ into $\omega\rho^-$ were found. The resonance in the $b_1(1235)\pi$ wave with exotic quantum numbers $J^{PC} = 1^{-+}$ at $M \approx 1.6 GeV$ is observed and the simultaneous analysis of the 1^{-+} wave in the $b_1(1235)\pi$, $\eta'\pi$ and $\rho\pi$ final states is presented.

1 Introduction

We present the results of the PWA of $\pi^+\pi^-\pi^-$, produced in reaction:

$$\pi^- Be \rightarrow \pi^+\pi^-\pi^- Be \quad (1)$$

and $\omega\pi^-\pi^0$ system, produced in:

$$\pi^- Be \rightarrow \omega\pi^-\pi^0 Be, \quad \omega \rightarrow \pi^+\pi^-\pi^0 \quad (2)$$

Our previous results of the analysis of reaction (1) were published in ^{1, 2, 3)}, and of reaction (2) were partially reported at the conferences ^{4), 5)}. The measurements




Figure 1: Main $J^{PC} = 1^{++}$ waves for $t' < 0.06 \text{ GeV}^2$ (a-c) and $0.06 < t' < 0.7 \text{ GeV}^2$ (d-f)

were carried out using VES spectrometer exposed by the 37 GeV momentum π^- beam. The description of the setup can be found in ⁶).

2 Results of the $\pi^+\pi^-\pi^-$ -system PWA.

The selection criteria for reaction (1) and the description of the PWA procedure can be found in ³). The relativistic covariant helicity formalism ⁷) is used to construct the amplitudes and the positively definite density matrix of the full rank. The largest waves of the $J^P = 0^-, 1^+, 2^-$ channels are decoupled from the other waves with the same $J^P M^\eta$ and are free to interfere with each other. The $6 \cdot 10^6$ events with $|t'| < 0.06 \text{ GeV}^2$ ¹ and $2 \cdot 10^6$ with $0.06 < |t'| < 0.7 \text{ GeV}^2$ are analyzed separately.

We present the main features of the most significant waves in the high 3π mass region.

$J^P M^\eta = 1^+ 0^+$. A peak in the $1^+ 0^+ D\rho$ wave (Fig.1(b)) and the shoulder

¹ $t' = t - t_{min}$, where t - momentum transfer from the beam to the final state squared, t_{min} -its minimum value.




Figure 2: Main $J^{PC} = 2^{-+}$ waves for $t' < 0.06 \text{ GeV}^2$

in the $1^+0^+S\rho$ wave (Fig.1(a), (d)) are observed at $M \approx 1.7 \text{ GeV}$ and are considered to be the $a'_1(1700)$ decay into the $\rho\pi$. The peak was fitted with the coherent sum of the Breit-Wigner resonance and the exponential background. The fit results in the following $a'_1(1700)$ parameters:

$$M = 1.80 \pm 0.05 \text{ GeV} \quad \Gamma = 0.23^{+0.10}_{-0.03} \text{ GeV};$$

where the errors are dominated by systematics. The $a'_1(1700)$ branching ratios into the D-wave $\rho\pi$ and P-wave $f_2(1270)\pi$ are found to be:

$$\frac{Br(a'_1(1700) \rightarrow (\rho\pi)_D)}{Br(a'_1(1700) \rightarrow (\rho\pi)_S)} < 0.35 \quad \frac{Br(a'_1(1700) \rightarrow f_2\pi)}{Br(a'_1(1700) \rightarrow (\rho\pi)_S)} < 0.23 \text{ at 95\% CL}$$

$J^P M^\eta = 2^-0^+$. A complicated wave behaviour for low t' shown in Fig.2 is described by the interplay of the $\pi_2(1670)$ and $\pi'_2(2100)$ states ^{3, 8}. A peak at $M \approx 1.7 \text{ GeV}$ in the F-wave $\rho\pi$ wave intensity is observed.

$J^P M^\eta = 3^+0^+$. A resonance-like signal is observed near $M \approx 1.8 \text{ GeV}$ in the $\rho\pi$ and $f_2\pi$ waves, see Fig.3. The phase of the $3^+0^+D\rho$ as related to the $2^-0^+Sf_2$ is shown in Fig.3(c) and is in accordance with the expectations for a resonance. The




Figure 3: The $3^+0^+D\rho$ (a), $3^+0^+Pf_2$ (b) wave intensities, the phase difference $\phi(3^+D\rho) - \phi(2^-Sf_2)$ (c) for $0.06 < t' < 0.7 \text{ GeV}^2$; the $3^+0^+D\rho$ (d), $3^+0^+Pf_2$ (e) and $3^+0^+S\rho_3$ (f) wave intensities for $t' < 0.06 \text{ GeV}^2$

simultaneous fit of the $f_2\pi$ intensity with the relativistic Breit-Wigner function and the $\rho\pi$ intensity with the incoherent sum of the Breit-Wigner and the Chebyshev polynomial background results in the following parameters of the resonance:

$$M = 1.86 \pm 0.02 \text{ GeV}, \quad \Gamma = 0.54 \pm 0.03 \text{ GeV}.$$

The relative probability of decay into the $f_2(1270)\pi$ and $\rho(770)\pi$ is as follows:

$$\frac{Br(a_3 \rightarrow f_2(1270)\pi)}{Br(a_3 \rightarrow \rho\pi)} = 0.5 \pm 0.1.$$

$J^P M^\pi = 4^+1^+$. A bump near $M \approx 2 \text{ GeV}$ is found in the $4^+1^+G\rho$ and $4^+1^+Ff_2$ waves produced at high t' (Fig.3). The phase of the $4^+1^+G\rho$ relative to the $\pi_2(1670)$ is in accordance with the expectation for the resonance. The simultaneous fit of the $f_2\pi$ intensity with the relativistic Breit-Wigner function and the $\rho\pi$ intensity with the incoherent sum of the Breit-Wigner and the Chebyshev polynomial background results in the following parameters of the resonance:

$$M = 1.95 \pm 0.02 \text{ GeV}, \quad \Gamma = 0.34 \pm 0.10 \text{ GeV}.$$




Figure 4: The $4^+1^+D\rho$ (a) and $4^+1^+Ff_2$ (b) wave intensities and the phase difference $\phi(4^+G\rho) - \phi(2^-Sf_2)$ for $0.06 < t' < 0.7 \text{ GeV}^2$.




Figure 5: The $J^P M^\eta = 1^-1^+S\rho$ wave intensity for $t' < 0.06 \text{ GeV}^2$ (a) and for $0.06 < t' < 0.7 \text{ GeV}^2$ (b).

We identify this object with the $a_4(2040)$, having the following branching ratio:

$$\frac{Br(a_4(2040) \rightarrow f_2(1270)\pi)}{Br(a_4(2040) \rightarrow \rho\pi)} = 0.5 \pm 0.2.$$

$J^P M^\eta = 1^-1^+$. The observation of a signal in the $1^-1^+P(\rho)$ wave with $M = 1.62 \pm 0.02 \text{ GeV}$, $\Gamma = 0.24 \pm 0.05 \text{ GeV}$ was previously reported as preliminary by the VES ⁹⁾. Later, the observation of the state with $M = 1593 \pm 8_{-47}^{+20} \text{ MeV}$, $\Gamma = 168 \pm 20_{-12}^{+150} \text{ MeV}$ was declared by the E852 ¹⁰⁾. It was shown in ¹¹⁾ the model dependence of the signal behaviour. We do not observe such a narrow signal (Fig. 5) in the fit results with the applied PWA model. However there appears a peak at $M \approx 1.6 \text{ GeV}$ with $\Gamma \approx 0.3 \text{ GeV}$ in the 1^-1^+ wave intensity, described by the leading term in the expansion of the density matrix in terms of the eigenvalues

(shown in Fig.7).




Figure 6: Wave intensities of: a) $0^-0^+P1(\rho)$, b) $2^-0^+P2(\rho)$ with $-t' < 0.08 \text{ GeV}^2$., Wave intensities of: c) $3^+0^+S(\rho_3)$, d) $2^+1^+S2(\rho)$, e) $4^+1^+D2(\rho)$, f) $1^-1^+S(b_1)$.

3 The results of the $\omega\pi^-\pi^0$ -system PWA

The selection criteria for reaction (2) and the description of the PWA procedure can be found in ¹²⁾. The results of the PWA are presented in Fig. 6.

$J^P M^\pi = 0^-0^+$. A peak in the region of 1.74 GeV with the flat background dominates in the wave $0^-0^+P(\rho)$ at low t' (Fig. 6(a)). The phase of this wave relative to the smooth $1^+0^+P(b_1)$ wave has the resonant behaviour. The resonance parameters were determined by the fit of the wave intensity with the incoherent sum of a relativistic Breit-Wigner function and a cubic polynomial and are as follows: the mass $M = 1.737 \pm 0.005 \pm 0.015 \text{ GeV}$ and the width $\Gamma = 0.259 \pm 0.019 \pm 0.06 \text{ GeV}$.

$J^P M^\pi = 2^-0^+$. A clear peak is observed at $M \sim 1.67 \text{ GeV}$ with $\Gamma \sim 0.2 \text{ GeV}$ (Fig. 6(b)). The resonant phase behaviour of the $2^-0^+P1(\rho)$ and $2^-0^+P2(\rho)$ waves relative to the $1^+0^+P(b_1)$ wave is observed. The resonance parameters of the $2^-0^+P2(\rho)$ peak were estimated in the same way as for $\pi(1740)$: $M = 1.687 \pm$

$0.009 \pm 0.015 \text{ GeV}$ and $\Gamma = 0.168 \pm 0.043 \pm 0.053 \text{ GeV}$. We identify this phenomenon with the decay of the $\pi_2(1670)$ into $\omega\rho$. The partial branching ratio was found by normalization to the decay $\pi_2(1670) \rightarrow f_2(1270)\pi^{-3})$, observed in the current experiment decay:

$$Br(\pi_2(1670)^- \rightarrow \omega\rho^-) = 0.027 \pm 0.004 \pm 0.01$$

The limits of the $\pi_2(1670)$ decay branching ratios are set at the 2σ confidence level:

$$Br(\pi_2(1670) \rightarrow \rho_1(1450)\pi) < 0.0036, \quad Br(\pi_2(1670) \rightarrow b_1\pi) < 0.0019.$$

$J^P M^\eta = 3^+0^+$. A peak at $M_{5\pi} \sim 2 \text{ GeV}$ and $\Gamma \sim 0.35 \text{ GeV}$ is clearly seen in the $3^+0^+S(\rho_3(1690))$ wave for events with low t' (Fig. 6(c)). However the resonant phase motion was not found. Such wave behaviour can be attributed to the Deck effect process¹³⁾.

$J^P M^\eta = 2^+1^+$. The intensive production and decay of the $a_2(1320)$ is the main process at low masses (Fig. 6(d)). The decay probability was found to be $Br(a_2(1320) \rightarrow \omega\pi^-\pi^0) = (5 \pm 1)\%$. We define the $a_2(1320)$ partial width as that of a Breit-Wigner function with the S-wave $\omega\pi^-\pi^0$ background. The nature of the 1.7 GeV mass structure is unknown. There may be another resonance or the opening of the $\omega\rho$ channel¹⁴⁾. We can not make preference to a particular hypothesis.

$J^P M^\eta = 4^+1^+$. The signal at $M \approx 2 \text{ GeV}$ (Fig. 6(e)) with the resonant phase behaviour can be identified as the $a_4(2040)$. The $a_4(2040)$ parameters are estimated by the fit with the incoherent sum of a D -wave relativistic Breit-Wigner function and a polynomial background to be: $M = 1.944 \pm 0.008 \pm 0.050 \text{ GeV}$ and $\Gamma = 0.324 \pm 0.026 \pm 0.075 \text{ GeV}$. The t' -dependence of the $a_4(2040)$ is identical to that of the $a_2(1320)$.

$J^P M^\eta = 1^-1^+$. The intensity of the $1^-1^+S(b_1)$ wave for events with high t' shows a wide bump with maximum at $M \sim 1.6 \div 1.7 \text{ GeV}$. The highest intensity of this wave does not exceed 15% of the $2^+1^+S2(\rho)$ wave intensity (Fig. 6(f)). The $\omega\rho$ P-waves are included in turn along with the $b_1\pi$ wave. Their intensity distribution differs in form from that of the $b_1\pi$ wave and their inclusion in the fit do not influence on the $b_1\pi$ wave intensity behaviour.

4 The $J^P M^\eta = 1^-1^+$ wave analysis

The combined analysis of the $J^P M^\eta = 1^-1^+ b_1\pi$ and the $J^P M^\eta = 2^+1^+ \omega\rho$ was carried out in order to understand the nature of the $b_1\pi$ wave. The results of the $\omega\pi^-\pi^0$ system PWA were used for this analysis. The diagonal elements and the real




Figure 7: Intensities of the $J^P M^\eta = 1^- 1^+$ waves in the channels: $b_1(1235)\pi$, $\eta'\pi$ and $\rho\pi$.

and imaginary parts of the non-diagonal element of the ρ -matrix corresponding to the $1^- 1^+$ $b_1\pi$ and $2^+ 1^+$ $\omega\rho$ waves with high t' were simultaneously fitted. Fit results were used to predict the coherence parameter for cross checking. The $b_1\pi$ amplitude was saturated by the Breit-Wigner resonance and the coherent background. The $\omega\rho^-$ amplitude was saturated by the $a_2(1320)$ -meson, a background and an a'_2 state has been also tried. The results of the fits with various ways of the $\omega\rho^-$ wave construction point out to the resonance nature of the $b_1\pi$ signal. The range of the parameters variation is large due to the freedom in the $2^+ 1^+$ state model.

The signal in the $J^P M^\eta = 1^- 1^+$ wave of the $\eta'\pi^-$ -system with close parameters was observed earlier¹⁵⁾. The simultaneous fit of the $b_1\pi$ and $\eta'\pi$ intensities with incoherent sum of a Breit-Wigner resonance and a background in each channel was carried out (Fig. 7). The fit results in the following parameters:

$$M = 1.58 \pm 0.03 \text{ GeV}, \quad \Gamma = 0.30 \pm 0.03 \text{ GeV}.$$

The form of the signal in the $1^- 1^+$ $\rho\pi$ wave is close to the Breit-Wigner function with these parameters. A fit of all three channels results in the resonance parameters changed within errors. All these facts indicate to the existence of a wide resonance $\Gamma = 0.29 \pm 0.03 \text{ GeV}$ with the mass $M = 1.61 \pm 0.02 \text{ GeV}$ and the relative branching ratio: $Br(b_1\pi) : Br(\eta'\pi) : Br(\rho\pi) = 1 : 1.0 \pm 0.3 : 1.6 \pm 0.4$.

5 Conclusions

The PWA of the reaction $\pi^- Be \rightarrow \pi^+ \pi^- \pi^- Be$ and $\pi^- Be \rightarrow \pi^+ 2\pi^- 2\pi^0 Be$ was performed.

The mass and the width of the resonance structure in the $J^P M^\eta = 0^- 0^+$ $\omega\rho^-$ wave differs from that for $\pi(1800)$. There may exist two objects of the different nature: a hybrid $\pi(1800)$ and a 3^1S_0 $q\bar{q}$ state decaying into $\omega\rho$.

The indication of the existence of the resonance a'_1 mostly decaying to the $\rho\pi$ in the S-wave is found.

The $\pi_2(1670)$ decays into the $\omega\rho$ and $\rho\pi$ in the F-wave are found.

The $a_2(1320)^- \rightarrow \omega\pi^-\pi^0$ decay and a wide bump of unknown nature at $M \approx 1.7 GeV$ are observed in the $J^P M^\eta = 2^+ 1^+$ wave. The a_3 and $a_4(2040)$ decays to the $\rho\pi$ and $f_2\pi$ are observed with the following relative branching ratio of the $a_4(2040)$ decays: $Br(f_2(1270)\pi) : Br(\rho\pi) : Br(\omega\rho) = 0.5 \pm 0.2 : 1 : 1.5 \pm 0.4$

The preliminary results of the 1^{-+} wave analysis point out to the existence of the resonance with exotic quantum numbers, which are forbidden for $q\bar{q}$ states, the mass $M = 1.61 \pm 0.02 GeV$, the width $\Gamma = 0.29 \pm 0.02 GeV$ and the following relative branching ratio: $Br(b_1\pi) : Br(\eta'\pi) : Br(\rho\pi) = 1 : 1.0 \pm 0.3 : 1.6 \pm 0.4$.

References

1. G. Beladidze *et al.*, Zeit. Phys. **C54** (1992), 367.
2. A. Zaitsev, "Recent results from VES detector at IHEP," Proc. of the IVth Int. Conf. on Hadron Spectroscopy (HADRON 91), 1991, 51.
3. D.I. Amelin *et al.*, Phys. Lett. **B356** (1995), 595.
4. D. Ryabchikov *et al.*, Proc. of the 6th Int. Conf. on Hadron Spectroscopy, (1995) 192.
5. A. Zaitsev *et al.*, Proc. of Int. Conf. on High Energy Physics, Warsaw, Poland, Jul. 1996
6. S. Bitjukov *et al.*, Phys. Lett. **B268** (1991), 137.
7. S.U.Chung, Phys. Rev. **D48** (1993), 1225;
F.Filippini *et al.*, Phys. Rev. **D51** (1995), 2247.
8. C. Daum *et al.*, Phys. Lett. **B89** (1980), 285.

9. Yu. Gouz *et al.*, Proc. of the XXVI International Conference on High Energy Physics, **1** 1992, 572.
10. A. Ostrovidov *et al.*, AIP Conf. Proc. 432, "Hadron Spectroscopy", Eds.S.-U. Chung and H. J. Willutzki, (Amer. Inst. Phys., New York, 1998), 263.
11. A. Zaitsev *et al.*, AIP Conf. Proc. 432, "Hadron Spectroscopy", Eds.S.-U. Chung and H. J. Willutzki, (Amer. Inst. Phys., New York, 1998), 461.
12. D. Amelin *et al.*, "Study of the reaction $\pi^- Be \rightarrow \pi^+ 2\pi^- 2\pi^0 Be$ ", hep-ex/9810013
13. R.T.Deck, Phys. Rev. Lett. **13** (1964), 169.
14. K.M.Watson. Phys. Rev. **88** (1952), 1163;
A.B.Migdal. Soviet Physics-JETP, **1** N. 1 (1955), 2.
15. G. Beladidze *et al.*, Phys. Lett. **B313** (1993), 276.